

An extension of the HarishChandra-Itzykson-Zuber integral

E. Brézin^{a)} and S. Hikami^{b)}^{a)} Laboratoire de Physique Théorique, Ecole Normale Supérieure
24 rue Lhomond 75231, Paris Cedex 05, France¹^{b)} Department of Basic Sciences,
University of Tokyo, Meguro-ku, Komaba, Tokyo 153, Japan**Abstract**

The HarishChandra-Itzykson-Zuber integral over the unitary group $U(k)$ ($\beta = 2$) is present in numerous problems involving Hermitian random matrices. It is well known that the result is semi-classically exact. This simple result does not extend to other symmetry groups, such as the symplectic or orthogonal groups. In this article the analysis of this integral is extended first to the symplectic group $Sp(k)$ ($\beta=4$). There the semi-classical approximation has to be corrected by a WKB expansion. It turns out that this expansion stops after a finite number of terms ; in other words the WKB approximation is corrected by a polynomial in the appropriate variables. The analysis is based upon new solutions to the heat kernel differential equation. We have also investigated arbitrary values of the parameter β , which characterizes the symmetry group. Closed formulae are derived for arbitrary β and $k = 3$, and also for large β and arbitrary k .

¹ Unité Mixte de Recherche 8549 du Centre National de la Recherche Scientifique et de l'École Normale Supérieure.

1 Introduction

The HarishChandra and Itzykson-Zuber (HIZ) integral [1, 2, 3] for the unitary group U_k is defined by

$$I = \int dU \exp i \text{tr} U^\dagger X U \Lambda \quad (1)$$

where X and Λ are diagonal $k \times k$ Hermitian matrices. It is well known that this integral is given, up to some normalization of the measure, by

$$I = \frac{\det_{1 \leq i, j \leq k} \exp(i x_i \lambda_j)}{\Delta(X) \Delta(\Lambda)} \quad (2)$$

where $\Delta(X)$ and $\Delta(\Lambda)$ are the Vandermonde determinants of the eigenvalues of X and Λ , ($\Delta(X) = \prod_{i < j} (x_i - x_j)$). This HIZ formula may be easily derived by considering the Laplacian operator [2] [9]

$$L = -\frac{\partial^2}{\partial X_{ij}^2}. \quad (3)$$

Its eigenfunctions are plane waves

$$L e^{i \text{tr} \Lambda X} = (\text{tr} \Lambda^2) e^{i \text{tr} \Lambda X}. \quad (4)$$

One can construct a unitary invariant eigenfunction of L , for the same eigenfunction $\text{tr} \Lambda^2$, by the superposition

$$I = \int dU e^{i \text{tr} \Lambda U X U^\dagger}, \quad (5)$$

which is nothing but the HIZ integral. The integral I being unitary invariant, is a function of the k eigenvalues x_i of X . The one dimensional quantum mechanics of the eigenvalues, for unitary invariant eigenstates, is then easily solved in terms of free fermions, leading to the unitary HIZ formula.

The same considerations hold for the three ensembles $\beta = 1, 2$ and 4 , corresponding to the orthogonal, unitary and symplectic ensembles, with

$$I = \int e^{i \text{tr} \Lambda g X g^{-1}} dg. \quad (6)$$

The Laplacian acting on invariant states, may be expressed in terms of a differential operator on the eigenvalues x_i :

$$[\sum_{i=1}^k \frac{\partial^2}{\partial x_i^2} + \beta \sum_{i=1, (i \neq j)}^k \frac{1}{x_i - x_j} \frac{\partial}{\partial x_i}] I = -\epsilon I , \quad (7)$$

with the eigenvalue ϵ

$$\epsilon = \sum_{i=1}^k \lambda_i^2 \quad (8)$$

Note that the integral I is manifestly symmetric under interchange of the matrices Λ and X , but the procedure is dissymmetric. The solutions will of course restore this property, which is not obvious if one considers the equation (7) alone.

The x -dependent eigenfunctions of this Schrödinger operator have a scalar product given by the measure

$$\langle \varphi_1 | \varphi_2 \rangle = \int dx_1 \cdots dx_k |\Delta(x_1 \cdots x_k)|^\beta \varphi_1^*(x_1 \cdots x_k) \varphi_2(x_1 \cdots x_k) \quad (9)$$

The measure becomes trivial if one multiplies the wave function by $|\Delta|^{\beta/2}$. Thus if, for some given ordering of the x_i 's, one changes $I(x)$ to

$$\psi(x_1 \cdots x_k) = |\Delta(x_1 \cdots x_k)|^{\beta/2} I(x_1 \cdots x_k), \quad (10)$$

one obtains the Hamiltonian,

$$[\sum_{i=1}^k \frac{\partial^2}{\partial x_i^2} - \beta(\frac{\beta}{2} - 1) \sum_{i < j} \frac{1}{(x_i - x_j)^2}] \psi = -\epsilon \psi. \quad (11)$$

The relation between matrix quantum mechanics and many-body problems with $1/r^2$ pair potentials was already present in [7] and it has also be used for the study of Selberg integrals by Forrester [8]. This Schrödinger equation is a simple Calogero-Moser model.

Although the solutions of this many-body problem are known, they are of little use for our purpose, since it is already quite involved to recover from the formal expression of the solution for arbitrary β , the simple unitary result of

HIZ and it looks formidable to extend it to other values of β [10]. For $\beta = 2$, the solution is again given by plane waves in the x_i and (taking into account the symmetry under permutations of I), one obtains the HIZ formula. For $\beta = 4$, the problem is much less trivial, but it leads to explicit solutions for finite values of k . Indeed the problem turns out to possess simple solutions of the form of symmetrized sums of plane waves multiplied by polynomials in the variables $1/(x_i - x_j)(\lambda_i - \lambda_j)$, providing therefore a complete explicit solution for the $Sp(k)$ group integration.

For general k , the solution of (11) is of the form

$$\psi_0 = e^{i(\lambda_1 x_1 + \dots + \lambda_k x_k)} \chi \quad (12)$$

where χ satisfies

$$\left[\sum_{i=1}^k \frac{\partial^2}{\partial x_i^2} + 2i \sum_{i=1}^k \lambda_i \frac{\partial}{\partial x_i} - y \sum_{i < j} \frac{1}{(x_i - x_j)^2} \right] \chi = 0 \quad (13)$$

in which

$$y = \beta \left(\frac{\beta}{2} - 1 \right). \quad (14)$$

For $k=2$, it is immediate to verify that there is a solution, function of the single variable $\tau = (\lambda_1 - \lambda_2)(x_1 - x_2)$, whose asymptotic expansion for large τ is

$$\begin{aligned} \chi &= 1 + \sum_{n=1}^{\infty} (-i/\tau)^n \prod_1^n \left(p - 1 - \frac{y}{2p} \right) \\ &= I_\nu(z) \sqrt{2\pi z} e^{-z} \end{aligned} \quad (15)$$

in which the Bessel function $I_\nu(z)$ is defined with $\nu^2 = (y + 1/2)/4$ and $z = i\tau/4$.

When $\beta = 4$, the operator $\sum_{i=1}^k \frac{\partial^2}{\partial x_i^2} - \sum_{i < j} \frac{4}{(x_i - x_j)^2}$ annihilates the function $\Delta^{-1}(x_1, \dots, x_k)$. Consequently the solution of (13) may be written

$$\chi(x_1 \dots x_k) = \frac{f(x_1 \dots x_k)}{\Delta(x_1 \dots x_k)} \quad (16)$$

in which $f(x_1 \cdots x_k)$ is a polynomial of degree $k(k-1)/2$ in the x_a 's. Defining

$$\tau_{ij} = (\lambda_i - \lambda_j)(x_i - x_j) \quad (17)$$

one finds for $k = 3$

$$\begin{aligned} \chi = & 1 + 2i\left(\frac{1}{\tau_{12}} + \frac{1}{\tau_{23}} + \frac{1}{\tau_{31}}\right) - 4\left(\frac{1}{\tau_{12}\tau_{23}} + \frac{1}{\tau_{23}\tau_{31}} + \frac{1}{\tau_{31}\tau_{12}}\right) \\ & - 12i\frac{1}{\tau_{12}\tau_{23}\tau_{31}}. \end{aligned} \quad (18)$$

Remaining still with $\beta = 4$, one may factor out the Vandermonde determinant from χ as (16), and one obtains a differential equation for f ,

$$\sum_{i=1}^k \frac{\partial^2}{\partial x_i^2} f + 2\left(\frac{\beta}{2} - 1\right) \sum_{i=1}^k \left(\frac{\partial f}{\partial x_i} + i\lambda_i f\right) \left(\Delta \frac{\partial}{\partial x_i} \frac{1}{\Delta}\right) + 2i \sum_{i=1}^k \lambda_i \frac{\partial f}{\partial x_i} = 0 \quad (19)$$

The solution of this equation is obtained by expanding in powers of the λ_i 's, but the expansion ends with the highest degree, the Vandermonde product of degree six $\Delta(\lambda_1 \cdots \lambda_4)$. Using the notation of (17), $\tau_{ij} = (x_i - x_j)(\lambda_i - \lambda_j)$, we obtain

$$f = 1 - \frac{i}{4}(\tau_{12} + \tau_{13} + \tau_{14} + \tau_{23} + \tau_{24} + \tau_{34}) + O(\tau^2) \quad (20)$$

as shown in the appendix of [4]. In the next section, we consider the symplectic case ($\beta = 4$) for larger values of k , and we will clarify the meaning of the coefficients which appear in the expansion of χ .

2 Decomposition into complete graphs in the symplectic case

Let us discuss now the asymptotic expansion of χ for $\beta = 4$, i.e. for the symplectic group integration of the HIZ integral. Such integrals appear in various problems ; for instance one encounters it in the calculation of the average of products of characteristic polynomials of random matrices in the

Gaussian orthogonal ensemble $\langle \prod_{i=1}^k \det(\lambda_i - X) \rangle$. Indeed we have shown in a previous work [4] that this average may be expressed as an integral over a $k \times k$ quaternion matrix ; this representation exhibits a duality between the GOE average and a symplectic HIZ integral. Thus this requires to compute the HIZ integral for $\beta = 4$, as discussed in [4] , in which the expressions were given up to $k = 4$. We extend here the construction of the polynomial solutions to arbitrary values of k .

As discussed before, the expansion of the function χ in inverse powers of the variables $1/\tau_{ij}$, defined in (17), terminates with a term proportional to $\prod_{i < j} \frac{1}{\tau_{ij}}$, of degree $k(k-1)/2$. The monomials in the variables $1/\tau_{ij}$ may be represented graphically : one marks k points, and a line between the points i and j will represent a factor $\frac{1}{\tau_{ij}}$ in a given monomial. A *complete* graph is a graph in which all the pairs of points are connected by a line. For instance, a triangle ($k=3$) and a tetrahedron ($k=4$) are complete graphs. The k -point complete graph has $k(k-1)/2$ lines. A complete graph of k points receives a factor C_k in which

$$C_k = \prod_{l=1}^k l! \quad (21)$$

This number is related to the integral,

$$C_k = \frac{1}{(2\pi)^{k/2}} \int \prod_{i < j} (z_i - z_j)^2 e^{-\frac{1}{2} \sum_i z_i^2} \prod_i dz_i \quad (22)$$

The last term of highest degree in χ , is equal to

$$i^k C_k \prod_{i < j} \frac{1}{\tau_{ij}} \quad (23)$$

All the terms of the expansion of χ have coefficients which may be decomposed into combinations of factors C_n with $n \leq k$.

For instance , the $k=3$ solution is

$$\chi = 1 + i \frac{C_2 C_1}{C_0} (v_1 + v_2 + v_3) + i^2 \frac{(C_2)^2}{C_1} (v_1 v_2 + v_2 v_3 + v_1 v_3) + i^3 C_3 v_1 v_2 v_3 \quad (24)$$

where $v_3 = \frac{1}{\tau_{12}}$, $v_1 = \frac{1}{\tau_{23}}$, and $v_2 = \frac{1}{\tau_{31}}$. The graphs of the various terms in χ may all be analyzed as a decomposition into complete graphs. For instance, $v_1 v_2$ consists of two lines connecting the point number three to points one and two. The lines (2-3) and (3-1) are complete graphs of two points, hence the factor C_2^2 , with the point three counted twice, thus a division by (C_1) , equal to one. (One has to multiply further the result by a power of i raised to the number of lines).

When $k = 4$, by this rule of decomposition of the graphs into complete graphs, we find the coefficients of all the terms in the form

$$\begin{aligned}
\chi = & 1 + i \frac{C_2 C_1}{C_0} \sum_{i=1}^6 v_i \\
& + i^2 \frac{(C_2)^2}{C_1} (v_1 v_2 + v_2 v_3 + v_1 v_3 + v_1 v_4 + v_1 v_5 + v_2 v_5 + v_2 v_6 + v_3 v_6 \\
& + v_3 v_4 + v_4 v_5 + v_4 v_6 + v_5 v_6) \\
& + i^2 (C_2)^2 (v_1 v_6 + v_2 v_4 + v_3 v_5) \\
& + i^3 \frac{C_3 C_1}{C_0} (v_1 v_2 v_3 + v_1 v_4 v_5 + v_2 v_5 v_6 + v_3 v_4 v_6) \\
& + i^3 \frac{(C_2)^3}{(C_1)^2} (v_1 v_3 v_5 + v_1 v_3 v_6 + v_2 v_3 v_4 + v_3 v_4 v_5 + v_3 v_5 v_6 + v_1 v_5 v_6 + v_2 v_4 v_5 \\
& + v_2 v_3 v_5 + v_1 v_2 v_4 + v_1 v_4 v_6 + v_1 v_2 v_6 + v_2 v_4 v_6) \\
& + i^3 \frac{(C_2)^3}{C_1} (v_1 v_2 v_5 + v_1 v_3 v_4 + v_2 v_3 v_6 + v_4 v_5 v_6) \\
& + i^4 \frac{(C_2)^4 C_0}{(C_1)^4} (v_1 v_2 v_4 v_6 + v_1 v_3 v_5 v_6 + v_2 v_3 v_4 v_5) \\
& + i^4 \frac{C_3 C_2}{C_1} (v_2 v_4 v_5 v_6 + v_2 v_3 v_5 v_6 + v_1 v_2 v_5 v_6 + v_3 v_4 v_5 v_6 + v_1 v_3 v_4 v_6 + v_2 v_3 v_4 v_6 \\
& + v_1 v_2 v_3 v_5 + v_1 v_2 v_3 v_6 + v_1 v_2 v_3 v_4 + v_1 v_3 v_4 v_5 + v_1 v_2 v_4 v_5 + v_1 v_4 v_5 v_6) \\
& + i^5 \frac{(C_3)^2}{C_2} (v_2 v_3 v_4 v_5 v_6 + v_1 v_3 v_4 v_5 v_6 + v_1 v_2 v_4 v_5 v_6 + v_1 v_2 v_3 v_5 v_6 \\
& + v_1 v_2 v_3 v_4 v_6 + v_1 v_2 v_3 v_4 v_5) \\
& + i^6 C_4 v_1 v_2 v_3 v_4 v_5 v_6
\end{aligned} \tag{25}$$

where we have used $v_1 = 1/\tau_{12}$, $v_2 = 1/\tau_{23}$, $v_3 = 1/\tau_{13}$, $v_4 = 1/\tau_{14}$, $v_5 =$

$1/\tau_{24}, v_6 = 1/\tau_{34}$. Again the graph of the last term $v_1 v_2 v_3 v_4 v_5 v_6$ is a complete graph of 4-points. Substituting in this formula the values $C_n = \prod_{j=1}^n j!$, one recovers precisely the $k = 4$ result, given in an earlier publication[4, 6].

The decomposition of graphs into complete components has certain general structures for arbitrary values of k . The term of highest degree is $i^{k(k-1)/2} C_k$; the next one, obtained by deleting one line, has a decomposition given by

$$\frac{(C_{k-1})^2}{C_{k-2}} \quad (26)$$

For $k=5$, this decomposition means that the graph is made of two tetrahedra sharing a triangular face; hence the division by C_3 to avoid double counting of this face. This explains the above factor (26). In the case $k=4$, this factor represents two triangles connected by one edge, and it reads $\frac{(C_3)^2}{C_2} = 72$.

When two lines are eliminated from the maximum complete graph C_k , we have two type graphs.

(i) The first case is the elimination of two lines emerging from the same point. In this case the decomposition is $C_{k-1} C_{k-2} / C_{k-3}$. For $k=4$, it is a triangle and one line connected at the vertex. For $k=5$, it is a tetrahedron connected to a triangle through one edge.

(ii) The second type consists of deleting two non consecutive lines. In this case, we have the decomposition of $(C_{k-2})^4 C_{k-4} / (C_{k-3})^4$. In the case $k=4$, it corresponds just to a square, four line connecting the four vertices of a square. For $k=5$, it is a graph of four triangles connected to each other by four edges.

If we divide this factor $(C_{k-2})^4 C_{k-4} / (C_{k-3})^4$ by C_k , as appeared in the expression for f in (20), it becomes $\frac{1}{18}, \frac{3}{80}, \frac{2}{75}$ for $k=4, 5$ and 6 , respectively. These values have been found by a direct evaluation of the solution of the differential equation (19). Thus we have, for $\beta = 4$ and for arbitrary values of k , an expression for χ , whose coefficients are given by the decomposition

of the complete graphs C_n .

The case $\beta = 4$ is thus easy to analyze for two reasons. First there are no "multiple" lines in a graph, meaning that every monomial in χ is of degree at most one in any variable $1/\tau_{ij}$. Then the factor $C_n = \prod j!$ for complete graphs has a simple structure. However, for $\beta \neq 4$, the number of lines between two points may be arbitrarily large. Then the contribution C_n of complete graphs has a more complicated expression. For values of β of the form

$$\beta = 2(m + 1) \quad (27)$$

a complete graph between two points consists of m lines joining these two points and

$$C_2 = \frac{y}{2}(-1 + \frac{y}{4})(-2 + \frac{y}{6}) \cdots (-(m - 1) + \frac{y}{2m}) \quad (28)$$

in which y has been defined in (14). The evaluation of C_3 is non trivial and we will discuss it in the following section. For β 's of the form (27) the previous graphical analysis of decomposition into complete subgraphs, may be extended to arbitrary k . However, even then, more complex rules for non-complete graphs are required. To understand this complexity, we consider the $k = 3$ case for arbitrary β in the next section.

3 $k=3$ for arbitrary β

Even for the simple $k = 3$ case, the HIZ integral is not known explicitly in the form of a WKB expansion. An analysis in terms of the solutions of the Calogero three-body problem has been performed in [10]. However, its outcome is far from our purpose since it is already quite involved to recover from there the unitary HIZ formula, and one does not see that for values of β such as four, the WKB expansion is a polynomial. Therefore we start again with the partial derivative equation (13) for $k = 3$, and expand the solution

as

$$\chi = \sum_{n,m,r=0} i^{n+m+r} \frac{C_{n,m,r}}{\tau_{12}^n \tau_{23}^m \tau_{31}^r}. \quad (29)$$

Note that it is not at all obvious that the equation (13) admits such a solution. Indeed, assuming that the solution is only a function of the τ 's, one finds easily that

$$[2i \sum_{i=1}^3 \lambda_i \frac{\partial}{\partial x_i} - \sum_{i < j} \frac{y}{(x_i - x_j)^2}] \chi = -(\lambda_1 - \lambda_2)^2 \tau_{12}^2 [2i \frac{\partial}{\partial \tau_{12}} + y] \chi + \text{perm.} \quad (30)$$

However the term involving the second derivative does not share this structure since

$$\begin{aligned} \sum_1^3 \frac{\partial^2 \chi}{\partial x_a^2} = & 2(\lambda_1 - \lambda_2)^2 [\tau_{12}^4 \frac{\partial^2 \chi}{\partial \tau_{12}^2} + \tau_{12}^3 \frac{\partial}{\partial \tau_{12}}] \\ & - 2(\lambda_1 - \lambda_2)(\lambda_3 - \lambda_1) \tau_{12}^2 \tau_{31}^2 \frac{\partial^2 \chi}{\partial \tau_{12} \partial \tau_{31}} + \text{perm} \end{aligned} \quad (31)$$

The last term, with cross-products in the λ_a 's, is a priori different. However there are, order by order, a set of identities which allow one to cast those cross-products into sums of squares. Those identities follow from $(x_1 - x_2) + (x_2 - x_3) + (x_3 - x_1) = 0$, leading to

$$(\lambda_1 - \lambda_2) \tau_{23} \tau_{31} + \text{perm} = 0. \quad (32)$$

From there one derives order by order the identities which are needed to cast the cross-products into squares. Let us quote the simplest

$$(\lambda_1 - \lambda_2)(\lambda_3 - \lambda_1) \tau_{23}^2 + \text{perm} = (\lambda_1 - \lambda_2)^2 \tau_{31} \tau_{23} + \text{perm.} \quad (33)$$

Order by order those identities allow us to satisfy the conditions on the expansion of χ , which make it a function of the τ variables. One finds easily that C_{n00} , which is associated to the diagram consisting of n lines between two points, is given by :

$$C_{n00} = \prod_{l=1}^n \left(-(l-1) + \frac{y}{2l} \right) \quad (34)$$

The generating function of those coefficients is the modified Bessel function, solution of the $k=2$ problem. Note that $C_{n,m,r}$ is symmetric under permutations of n, m and r ; for instance $C_{m,0,0} = C_{0,m,0}$.

When $r=0$, one finds that $C_{n,m,0}$ has a decomposition into a product of two factors,

$$C_{n,m,0} = C_{n,0,0} C_{m,0,0}. \quad (35)$$

For the full HIZ integral, triangle graphs, in which the three integers (n, m, r) of $C_{n,m,r}$ are non-zero, play an essential role.

From the differential equation (13), one can derive a recursion equation for the $C_{n,m,r}$. However a direct expansion of the solution of is extremely tedious and complicated at increasing orders, since one has to use more and more identities of the type (33), but of higher degree. One needs triangle identities. The expansion for the differential equation (19) has a simpler structure ; it gives a relation between $C_{n,m,r}$, with $n + m + r = P + 1$, in terms of $C_{n,m,r}$ with $n + m + r = P$; this relation is valid for a fixed given value of m . If one writes $C_{n,m,r}$ as a linear combination of $C_{n',m,r'}$, where $r' < r$, and $n' + r' = n + r - 1$, one obtains

$$C_{n,m,1} = C_{n,m,0} \left(nm + \frac{y}{2} \right) \quad (36)$$

$$C_{n,m,2} = C_{n,m,1} \left(\frac{m(n-1)}{2} - 1 + \frac{y}{4} \right) + C_{n+1,m,0} \frac{m}{2} (n+1) \quad (37)$$

$$\begin{aligned} C_{n,m,3} = & C_{n,m,2} \left(\frac{m}{3} (n-2) - 2 + \frac{y}{6} \right) + C_{n+1,m,1} \frac{mn}{3} \\ & + C_{n+2,m,0} \frac{m}{3} (n+2) \end{aligned} \quad (38)$$

For general r ($r > 3$), we have

$$\begin{aligned} C_{n,m,r} = & C_{n,m,r-1} \left(\frac{m(n-r+1)}{r} - (r-1) + \frac{y}{2r} \right) \\ & + C_{n,m,r-2} \frac{m}{r} (n-r+3) + C_{n,m,r-3} \frac{m}{r} (n-r+5) \end{aligned}$$

$$\begin{aligned}
& + \dots \\
& + C_{n+r-2,m,1} \frac{m}{r} (n+r-3) + C_{n+r-1,m,0} \frac{m}{r} (n+r-1). \quad (39)
\end{aligned}$$

The solution of these recursion equations guarantees automatically the symmetry under exchange between n, m and r : $C_{n,m,r} = C_{n,r,m} = C_{m,n,r} = C_{m,r,n} = C_{r,n,m} = C_{r,m,n}$.

From these recursion formulae, one may determine iteratively all the coefficients $C_{n,m,r}$. The resulting expressions are complicated for general y . For instance, even at low order, one finds

$$C_{2,2,2} = \left(\frac{y}{2}\right)^2 \left(-1 + \frac{y}{4}\right)^2 \left(-6 + \frac{3}{2}y + \frac{y^2}{8}\right) \quad (40)$$

$$C_{3,3,3} = \left(\frac{y}{2}\right)^2 \left(-1 + \frac{y}{4}\right)^2 \left(-2 + \frac{y}{6}\right)^2 \left(-51 - \frac{59}{8}y + \frac{37}{48}y^2 + \frac{1}{48}y^3\right) \quad (41)$$

The solution for $\beta = 4$ and $k = 3$, may be written in the compact form

$$\chi = e^\phi \quad (42)$$

$$\phi = 2i \left(\frac{1}{\tau_{12}} + \frac{1}{\tau_{23}} + \frac{1}{\tau_{13}} \right) - 4i \frac{1}{\tau_{12}\tau_{23}\tau_{13}} \quad (43)$$

in which it is understood that one expands it in powers of $1/\tau$, dropping any term involving powers of $1/\tau_{ij}$ greater than one. This corresponds to the absence of multiple bonds between two points in the graphical representation for $\beta = 4$. This result agrees with (18). This certainly suggests that ϕ ought to be written in terms of Grassmann variables. In the next section, we consider such expressions for general y in an expansion in $1/y$. This also provides consistency checks for the recursion equation (39).

4 Large y expansion

One can obtain an expansion in powers of $1/y$ if one returns to the differential equation (13), taking now as variables

$$u_a = x_a/y. \quad (44)$$

The function $\chi(u_1 \cdots u_k)$ is solution of

$$\left[\frac{1}{y} \sum_{a=1}^k \frac{\partial^2}{\partial u_a^2} + 2i \sum_{a=1}^k \lambda_a \frac{\partial}{\partial u_a} - \sum_{a < b} \frac{1}{(u_a - u_b)^2} \right] \chi = 0. \quad (45)$$

The leading term χ_0 is thus solution of

$$\left[2i \sum_{a=1}^k \lambda_a \frac{\partial}{\partial u_a} - \sum_{a < b} \frac{1}{(u_a - u_b)^2} \right] \chi_0 = 0. \quad (46)$$

We look for a solution which is a symmetric function of the variables

$$v_{a,b} = \frac{i}{(\lambda_a - \lambda_b)(u_a - u_b)}, \quad a < b \quad (47)$$

which thus satisfies

$$\sum_{a=1}^k \lambda_a \frac{\partial \chi_0}{\partial u_a} = i \sum_{1 \leq a < b \leq k} (\lambda_a - \lambda_b)^2 v_{ab}^2 \frac{\partial \chi_0}{\partial v_{ab}}. \quad (48)$$

The equation (46) takes then the simple form

$$\sum_{1 \leq a < b \leq k} (\lambda_a - \lambda_b)^2 v_{ab}^2 \left[2 \frac{\partial}{\partial v_{ab}} - 1 \right] \chi_0 = 0. \quad (49)$$

There is a trivial symmetric solution which satisfies the $k(k-1)/2$ equations

$$\left[2 \frac{\partial}{\partial v_{ab}} - 1 \right] \chi_0 = 0, \quad (50)$$

namely

$$\chi_0 = \exp \left(\frac{1}{2} \sum_{a < b} v_{ab} \right) = \exp \left(\frac{iy}{2} \sum_{a < b} \frac{1}{(\lambda_a - \lambda_b)(x_a - x_b)} \right). \quad (51)$$

One thus writes

$$\begin{aligned} \chi &= e^\phi \\ \phi &= \phi_0 + \frac{1}{y} \phi_1 + \frac{1}{y^2} \phi_2 + \cdots, \end{aligned} \quad (52)$$

with

$$\phi_0 = \frac{1}{2} \sum_{a < b} v_{ab}. \quad (53)$$

The next terms are solutions of

$$\sum_1^k \left(\left(\frac{\partial \phi_0}{\partial u_a} \right)^2 + \frac{\partial^2 \phi_0}{\partial u_a^2} \right) + 2i \sum_1^k \lambda_a \frac{\partial \phi_1}{\partial u_a} = 0, \quad (54)$$

$$\sum_1^k \left(2 \frac{\partial \phi_0}{\partial u_a} \frac{\partial \phi_1}{\partial u_a} + \frac{\partial^2 \phi_1}{\partial u_a^2} \right) + 2i \sum_1^k \lambda_a \frac{\partial \phi_2}{\partial u_a} = 0. \quad (55)$$

In order to make it clear that it is a non trivial property of these equations to have solutions which are symmetric polynomials in the variables v_{ab} , we shall examine these equations in some detail.

For simplicity of notations we shall limit ourselves to $k=3$, the generalization to arbitrary k being immediate. It is thus convenient to simplify slightly the notations and write

$$v_{12} = V_3, \quad v_{23} = V_1, \quad v_{31} = V_2 \quad (56)$$

Let us first consider (54) ; since the function ϕ_1 is a symmetric function of the V_i 's one finds easily, as in (48), that

$$\sum_1^3 \lambda_a \frac{\partial \phi_1}{\partial u_a} = i [(\lambda_2 - \lambda_3)^2 V_1^2 \frac{\partial}{\partial V_1} + \text{perm.}] \phi_1 \quad (57)$$

The explicit solution (53) allows one to write

$$\begin{aligned} \sum_1^3 \left(\left(\frac{\partial \phi_0}{\partial u_a} \right)^2 \right) &= -\frac{1}{2} [(\lambda_2 - \lambda_3)^2 V_1^4 + \text{perm.}] + \frac{1}{2} [(\lambda_1 - \lambda_2)(\lambda_3 - \lambda_1) V_2^2 V_3^2 + \text{perm.}] \\ \sum_1^3 \frac{\partial^2 \phi_0}{\partial u_a^2} &= -2 [(\lambda_2 - \lambda_3)^2 V_1^3 + \text{perm.}] \end{aligned} \quad (58)$$

The second term of the first line of (57) does not involve a priori $(\lambda_a - \lambda_b)^2$; however the variables V_i are not independent since they are defined in terms of the differences $(t_1 - t_2), (t_2 - t_3), (t_3 - t_1)$ whose sum is zero. Thus

$$V_2 V_3 (\lambda_1 - \lambda_2) (\lambda_1 - \lambda_3) + \text{perm.} = 0 \quad (59)$$

and one can check without difficulty that

$$V_2^2 V_3^2 (\lambda_1 - \lambda_2) (\lambda_1 - \lambda_3) + \text{perm.} = -V_1 V_2 V_3 [(\lambda_2 - \lambda_3)^2 V_1 + \text{perm.}]. \quad (60)$$

With the help of this identity the equation for ϕ_1 takes the simple form

$$(\lambda_2 - \lambda_3)^2 V_1^2 \left[\frac{\partial \phi_1}{\partial V_1} + \frac{1}{4} V_1^2 + V_1 - \frac{1}{4} V_2 V_3 \right] + \text{perm.} = 0. \quad (61)$$

Thanks to the identity (60) one can look for a solution of the three equations

$$\frac{\partial \phi_1}{\partial V_1} + \frac{1}{4} V_1^2 + V_1 - \frac{1}{4} V_2 V_3 = 0 \quad (62)$$

plus the other two deduced from the cyclic permutations. The system is trivially integrable and one finds

$$\phi_1 = \frac{1}{4} V_1 V_2 V_3 - \frac{1}{2} (V_1^2 + V_2^2 + V_3^2) - \frac{1}{12} (V_1^3 + V_2^3 + V_3^3). \quad (63)$$

For arbitrary k this generalizes to

$$\phi_1 = \frac{1}{4} \sum_{(abc)} v_{ab} v_{bc} v_{ca} - \frac{1}{2} \sum_{a < b} v_{ab}^2 - \frac{1}{12} \sum_{a < b} v_{ab}^3, \quad (64)$$

in which the sum (abc) runs over all distinct triplets of indices.

We have exposed this simple calculation in some detail, in order to make it clear that the success of the method relies on two successive facts :

- Identities such as (59, 60) which allows one to write the differential equation as three separate equations for $\partial \phi / \partial V_a$
- The integrability conditions of this system of three equations are satisfied.

At higher orders the technique is identical. It requires new identities of higher degree, such as

$$\begin{aligned} & V_2^2 V_3^2 (V_2^2 + V_3^2) (\lambda_1 - \lambda_2) (\lambda_1 - \lambda_3) + \text{perm.} \\ &= -(\lambda_2 - \lambda_3)^2 V_1 V_2 V_3 [V_1 (V_1^2 + V_2^2 + V_3^2) + \frac{1}{2} V_1 V_2 V_3] + \text{perm.} \end{aligned} \quad (65)$$

and many others ; thereby one obtains again remarkably an integrable system of three equations. At order $1/y^2$ the solution is

$$\begin{aligned}\phi_2 = & (V_1^3 + V_2^3 + V_3^3) + \frac{1}{2}(V_1^4 + V_2^4 + V_3^4) - \frac{1}{2}V_1V_2V_3(V_1 + V_2 + V_3) \\ & + \frac{1}{20}(V_1^5 + V_2^5 + V_3^5) - \frac{1}{8}V_1V_2V_3(V_1^2 + V_2^2 + V_3^2)\end{aligned}\quad (66)$$

5 A cubic identity

Let us consider the case of $k = 4$ for different values of β , assuming that it is an even integer. For the tetrahedron ($k=4$), the coefficients of the expansion of χ may be written as

$$\chi = \sum_{n,m,r,p,q,l=0}^j i^{n+m+r+p+q+l} \frac{C_{nmr;pql}}{\tau_{12}^n \tau_{13}^m \tau_{23}^r \tau_{24}^p \tau_{34}^q \tau_{14}^l} \quad (67)$$

where $j = \beta/2 - 1$. By imposing to satisfy the differential equation (19), one should determine those coefficients. For instance, in the case $\beta = 6$ (or $y = 12$), one finds $C_{222;222} = 8 \cdot 7 (6 \cdot 5)^2 (4 \cdot 3)^3$, $C_{222;221} = (2 \cdot 7) (6 \cdot 5)^2 (4 \cdot 3)^3$, $C_{222;211} = 4(6 \cdot 5)^2 (4 \cdot 3)^3$, $C_{222;220} = (C_{222;000})^2 / C_{200;000} = (6 \cdot 5)^2 (4 \cdot 3)^3$, $C_{222;000} = (6 \cdot 5)(4 \cdot 3)^2$, $C_{221;221} = 6(6 \cdot 5)^3 2^5$, $C_{nm2;pq1} = C_{nm2;pq0}(np + mq + \frac{y}{2})$, $C_{nmr;pq0} = C_{nmr;000}C_{pqr;000}/C_{r00;000}$, $C_{222;pq2} = C_{222;pq1}(\frac{p}{2} + \frac{q}{2} - 1 + \frac{y}{4})$. (The coefficient $C_{nmr;000}$ is identical to the C_{nmr} of the $k=3$ case (29). Whenever the graphs reduce to triangles, one recovers the coefficients of the $k=3$ problem).

However the above representation (67) is in fact somewhat ambiguous. Indeed there is an interesting cubic identity between the six variables

$$\tau_{ab} = (x_a - x_b)(\lambda_a - \lambda_b), \quad 1 \leq a < b \leq 4,$$

which holds for every set of x_a 's and λ_a 's, namely

$$\begin{aligned}& \tau_{12}^2 \tau_{34} + \tau_{13}^2 \tau_{24} + \tau_{14}^2 \tau_{23} + \tau_{23}^2 \tau_{14} + \tau_{24}^2 \tau_{13} + \tau_{34}^2 \tau_{12} \\ & - \tau_{12} \tau_{34} (\tau_{23} + \tau_{24} + \tau_{14} + \tau_{13}) - \tau_{13} \tau_{24} (\tau_{14} + \tau_{12} + \tau_{23} + \tau_{34})\end{aligned}$$

$$\begin{aligned}
& -\tau_{14}\tau_{23}(\tau_{12} + \tau_{24} + \tau_{34} + \tau_{13}) \\
& +\tau_{12}\tau_{24}\tau_{14} + \tau_{13}\tau_{14}\tau_{34} + \tau_{23}\tau_{24}\tau_{34} + \tau_{12}\tau_{23}\tau_{13} = 0.
\end{aligned} \tag{68}$$

Consequently, when one divides the l.h.s. of this identity by the Vandermonde squared $\Delta(x)^2\Delta(\lambda)^2$, one generates an identity for the terms of the expansion of χ whose coefficients are $C_{221;220}, C_{221;211}, C_{122;121}$. Thus there is an arbitrariness in writing χ in the form of the expansion (67), in view of this cubic identity. For arbitrary k , there are $(k-2)(k-3)/2$ similar cubic identities, and therefore only $(2k-3)$ independent τ_{ab} .

For other values of β such as $\beta = 8$ for instance, by imposing the differential equation (19), one finds that the values of $C_{332;331}, C_{332;322}, C_{323;322}$ are not uniquely determined, because again of this same cubic identity. One may need to eliminate this arbitrariness by choosing some arbitrary value for some coefficient, decide that is it zero for instance. This cubic identity carries the problem over to higher orders. Indeed, multiplying this same cubic identity, by some symmetric function of the τ 's, such as $\sum \tau_{ij}$, one obtains an identity of order τ^4 , which again requires some arbitrary choice in the expression for χ at next order.

In summary, we have obtained expressions for the HarishChandra-Itzykson-Zuber integration for $k \times k$ matrices, for group integrals characterized by the parameter β . Our investigation relies upon new solutions to the heat kernel differential equation. Other methods such as supersymmetric quantum mechanics, combinatorial group analysis, or algebraic geometry may be of help in solving the numerous open questions concerning this problem.

Acknowledgements

It is a pleasure to thank Dr. Alfaro for interesting remarks concerning the relevance of supersymmetric quantum mechanics in this problem. S.H. has benefited from a Grant-in-Aid for Scientific Research (B) by JSPS.

References

- [1] Harish-Chandra, Proc. Nat. Acad. Sci. **42**, 252(1956).
- [2] C. Itzykson and J.-B. Zuber, J. Math. Phys. **21**, 411 (1980).
- [3] J. J. Duistermaat and G. H. Heckman, Invent. Math.**69**, 259 (1982).
- [4] E. Brézin and S. Hikami, Commun. Math. Phys. **223**, 363 (2001).
- [5] E. Brézin and S. Hikami, Commun. Math. Phys. **214**, 111 (2000).
- [6] T. Guhr and H. Kohler, J. Math. Phys. **43**, 2707 (2002).
- [7] E. Brézin, C. Itzykson, G. Parisi and J.-B. Zuber, Comm. Math. Phys.**59**, 35 (1978).
- [8] P.J. Forrester, Nucl. Phys. **B388**, 671 (1992).
- [9] E. Brézin, "Two dimensional quantum gravity and random surfaces", p.37, edited by D. J. Gross T. Piran and S. Weinberg,(1992),World Scientific, Singapore.
- [10] G. Mahoux, M. L. Mehta and J.-M. Normand, "Random Matrices and Their Applications", MSRI Publications, **40**, 301 (2001).